SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN

www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

"Compassion Fatigue in Mental Health Nurses: Recognition and Prevention"

Ramandeep Kaur[1], Dr. Payal Sharma[2]
[1]PhD Research Scholar, [2]Research Supervisor
[1,2]Index Nursing College,
Malwanchal University, Indore, MP, India
Date of Publication: 24/09/2025
DOI 10.5281/zenodo.24465123

Abstract: Compassion is a fundamental component of mental health nursing, enabling nurses to establish therapeutic relationships, understand patients' emotional experiences and provide holistic, person-centred care. However, continuous exposure to psychological suffering, trauma, suicidal behaviour, self-harm, aggression, emotional distress and prolonged patient dependency may place mental health nurses at increased risk of compassion fatigue. Compassion fatigue is a work-related phenomenon commonly discussed in relation to secondary traumatic stress and occupational burnout. It can progressively affect nurses' emotional wellbeing, physical health, professional relationships, clinical judgement and capacity to provide compassionate care. Unlike ordinary occupational stress, compassion fatigue may involve emotional exhaustion associated with repeatedly witnessing or empathically engaging with the traumatic experiences of others. Mental health nurses working in psychiatric hospitals, rehabilitation centres, community mental health services, emergency psychiatric units and addiction treatment facilities may encounter distinctive occupational stressors that contribute to this phenomenon. This narrative review examines the concept, contributing factors, manifestations, consequences, assessment and prevention of compassion fatigue among mental health nurses. Particular attention is given to the emotional and psychological demands of psychiatric nursing, early recognition of warning signs, professional quality-of-life assessment, clinical supervision, reflective practice, mindfulness, resilience development and organisational interventions. The review also explores the responsibilities of nursing administrators, psychiatric nurse educators and healthcare institutions in establishing supportive working environments. Evidence suggests that individual self-care strategies alone are insufficient to prevent compassion fatigue when organisational stressors remain unaddressed. A comprehensive approach integrating individual wellbeing, supportive leadership, adequate staffing, professional supervision, peer support and institutional mental health programmes is essential. Recognising compassion fatigue as an important occupational health concern can help protect nurses' wellbeing, improve workforce retention and maintain the quality and continuity of mental healthcare.

Keywords: *Compassion fatigue, mental health nurses, psychiatric nursing, secondary traumatic stress, occupational burnout, professional quality of life, emotional exhaustion, vicarious trauma, resilience, clinical supervision, self-care, workplace wellbeing, nursing interventions, prevention strategies.*

1. Introduction

Mental health nursing is a specialised professional discipline that requires a combination of clinical competence, emotional intelligence, therapeutic communication, empathy, patience and interpersonal sensitivity. Mental health nurses work with individuals experiencing a wide range of psychological conditions, including schizophrenia, mood disorders, anxiety disorders, personality disorders, substance use disorders, trauma-related conditions and suicidal behaviour. Their responsibilities extend beyond medication administration and clinical observation to include emotional support, crisis intervention, therapeutic engagement, rehabilitation, family education and the promotion of recovery-oriented care.

Unlike several other healthcare settings, psychiatric nursing frequently involves prolonged therapeutic relationships with individuals experiencing persistent psychological distress, interpersonal difficulties, behavioural disturbances and complex psychosocial challenges. Nurses may repeatedly listen to traumatic experiences, witness emotional suffering, manage aggression, support individuals during suicidal crises and provide care to patients who have experienced abuse, neglect or social rejection. Although these experiences are central to mental health nursing, continuous exposure to suffering may gradually affect the emotional resources of healthcare professionals.

Compassion fatigue has therefore gained increasing attention as an occupational concern among nurses and other professionals

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

involved in caring for vulnerable populations. Figley described compassion fatigue in relation to the emotional and psychological consequences of helping individuals who have experienced traumatic events [1]. It is frequently discussed as a combination of secondary traumatic stress and burnout, although these concepts represent related but distinguishable experiences [2].

Mental health nurses may be particularly affected because therapeutic relationships require sustained emotional engagement. Repeated exposure to patients' distress, combined with workload pressures, staff shortages, workplace aggression and insufficient professional support, can gradually reduce emotional availability and professional satisfaction.

Marshman, Hansen and Munro conducted a systematic review specifically examining compassion fatigue among mental health nurses. Their review included 12 studies and identified varying levels of compassion fatigue, with organisational leadership, positive workplace culture, clinical supervision, reflection and individual wellbeing emerging as potentially protective factors [3]. These findings demonstrate that compassion fatigue should not be regarded merely as an individual weakness or inability to manage stress. Instead, it represents an important issue requiring attention at both professional and organisational levels.

Understanding compassion fatigue is essential for psychiatric nurses, nursing educators, clinical supervisors and healthcare administrators because early recognition and appropriate preventive measures can protect professional wellbeing while supporting safe, therapeutic and compassionate patient care.

2. Understanding Compassion Fatigue in Mental Health Nursing

Compassion fatigue refers to the emotional and psychological strain associated with repeatedly caring for individuals experiencing suffering, trauma or severe distress. It may develop when the emotional demands of caregiving exceed an individual's available coping resources over an extended period.

Figley identified compassion fatigue as a consequence of empathic engagement with individuals experiencing traumatic stress [1]. Stamm subsequently developed the Professional Quality of Life framework, which explains the positive and negative effects of helping others through three related dimensions: compassion satisfaction, burnout and secondary traumatic stress [4].

In mental health nursing, compassion fatigue may emerge gradually as nurses repeatedly encounter patients' traumatic experiences, emotional crises, suicidal behaviour, self-destructive actions and persistent psychological difficulties. Nurses may begin to experience emotional exhaustion, reduced empathy, irritability, helplessness and difficulty maintaining professional boundaries.

It is important to recognise that compassion fatigue is not an independent psychiatric diagnosis. It is a descriptive occupational phenomenon that overlaps with several recognised psychological and work-related stress responses. Its manifestations may vary according to personal characteristics, clinical responsibilities, working environments and available support systems.

2.1 Compassion Fatigue and Burnout

Burnout is generally associated with chronic workplace stress that has not been successfully managed. The World Health Organization recognises burnout in ICD-11 as an occupational phenomenon characterised by energy depletion or exhaustion, increased mental distance or cynicism towards one's job, and reduced professional efficacy [5].

In mental health nursing, burnout may develop because of excessive workload, inadequate staffing, long working hours, administrative responsibilities, insufficient resources and limited professional recognition. Nurses may experience persistent tiredness, frustration, reduced motivation and dissatisfaction with their professional roles.

Compassion fatigue and burnout can coexist, but they are not interchangeable. Burnout primarily reflects chronic occupational stress, whereas compassion fatigue places particular emphasis on the emotional consequences of caring for individuals experiencing suffering and trauma. Distinguishing these concepts is important because preventive interventions may need to address both workplace conditions and secondary traumatic stress.

2.2 Compassion Fatigue and Secondary Traumatic Stress

Secondary traumatic stress refers to trauma-related psychological reactions that may occur when an individual is indirectly exposed to another person's traumatic experiences. Mental health nurses may repeatedly hear descriptions of abuse, violence, sexual assault, childhood trauma, suicide attempts and other distressing events.

Such exposure may produce intrusive thoughts, emotional distress, avoidance, heightened alertness and sleep disturbances. Nurses may find themselves repeatedly thinking about patients' traumatic experiences even after leaving the workplace.

Secondary traumatic stress is particularly relevant in psychiatric settings because therapeutic communication frequently involves exploring painful personal experiences. Without adequate professional supervision, emotional processing and recovery opportunities, repeated exposure may contribute to compassion fatigue [2,3].

copyright©: Scientific Journal

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

2.3 Compassion Satisfaction

Compassion satisfaction represents the positive emotional experience associated with helping others and finding meaning in professional caregiving. Mental health nurses may experience satisfaction when patients demonstrate recovery, achieve rehabilitation goals, improve interpersonal functioning or regain independence.

Compassion satisfaction is an important protective dimension of professional quality of life. Positive therapeutic relationships, professional recognition, supportive colleagues, meaningful work and opportunities to observe patient progress can strengthen professional fulfilment [4].

A healthy professional environment should therefore aim not only to reduce compassion fatigue but also to enhance compassion satisfaction.

3. Factors Contributing to Compassion Fatigue Among Mental Health Nurses

Compassion fatigue develops through interactions between individual vulnerabilities, clinical exposure, professional responsibilities and organisational conditions. Understanding these contributing factors helps nursing professionals identify high-risk situations and develop appropriate prevention strategies.

3.1 Continuous Exposure to Psychological Suffering

Mental health nurses frequently interact with individuals experiencing severe emotional distress, hopelessness, fear, grief, trauma and interpersonal difficulties. Listening repeatedly to traumatic narratives and witnessing patients' psychological deterioration may create significant emotional demands.

For example, a psychiatric nurse caring for an adolescent with a history of abuse may repeatedly encounter intense emotional reactions, self-harming behaviour and family-related distress. Similarly, nurses working in suicide prevention services may experience emotional strain when supporting patients with recurrent suicidal crises. Repeated exposure without adequate opportunities for emotional recovery may contribute to secondary traumatic stress.

3.2 Management of Aggression and Workplace Violence

Aggression and violence represent important occupational challenges in psychiatric nursing. Nurses may encounter verbal threats, physical assaults, intimidation, property destruction and unpredictable behavioural disturbances.

Managing these situations requires continuous observation, emotional regulation, risk assessment and therapeutic communication. Repeated exposure to threatening situations may produce fear, hypervigilance, frustration and emotional exhaustion.

The experience of workplace violence can also affect nurses' confidence and perception of personal safety. Inadequate institutional responses following aggressive incidents may further increase psychological distress.

3.3 Heavy Workload and Staffing Shortages

Insufficient staffing, high patient-to-nurse ratios, extended working hours and inadequate opportunities for rest may intensify occupational stress. Mental health nurses must often balance direct patient care with documentation, medication administration, multidisciplinary meetings, family counselling and administrative responsibilities.

When workload demands consistently exceed available resources, nurses may have insufficient time to establish meaningful therapeutic relationships. This conflict between professional expectations and practical working conditions may contribute to frustration, emotional exhaustion and reduced professional satisfaction.

3.4 Therapeutic Relationship and Emotional Involvement

The therapeutic relationship is central to mental health nursing. Nurses are expected to demonstrate empathy, active listening, emotional availability and professional sensitivity while maintaining appropriate therapeutic boundaries.

However, excessive emotional involvement may make it difficult for nurses to separate professional responsibilities from personal feelings. Nurses may become excessively concerned about particular patients, feel responsible for outcomes beyond their control or experience guilt following unsuccessful interventions.

Conversely, repeated emotional distress may lead some nurses to distance themselves from patients as a protective response. Both excessive involvement and emotional withdrawal may negatively affect therapeutic relationships.

3.5 Patient Suicide, Self-Harm and Unexpected Death

Caring for patients who engage in repeated self-harm or suicidal behaviour can create considerable emotional strain. Patient suicide or unexpected death may be particularly distressing, especially when nurses have established strong therapeutic relationships with the individual.

Nurses may experience guilt, sadness, helplessness, self-doubt and repeated questioning of their clinical decisions. These experiences may become more difficult when post-incident support, debriefing or professional counselling is unavailable.

3.6 Ethical Dilemmas and Moral Distress

Mental health nurses frequently encounter ethical challenges involving patient autonomy, involuntary treatment, restrictive

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

interventions, confidentiality, patient rights and safety considerations.

Situations in which nurses believe they know the most therapeutic course of action but cannot implement it because of organisational restrictions or competing ethical responsibilities may produce moral distress.

Repeated exposure to ethically challenging situations without adequate professional discussion can contribute to emotional exhaustion and reduced job satisfaction.

3.7 Lack of Organisational Support

Inadequate managerial support, poor communication, workplace conflict, limited recognition and insufficient opportunities for professional development may increase vulnerability to compassion fatigue.

A workplace that discourages emotional disclosure or interprets psychological distress as professional incompetence may prevent nurses from seeking assistance. Conversely, supportive leadership and positive organisational cultures can strengthen professional resilience [3].

4. Recognition and Clinical Manifestations of Compassion Fatigue

Early identification is essential because compassion fatigue may develop gradually and can initially be mistaken for ordinary tiredness, temporary occupational stress or reduced motivation. Nurses and nursing administrators should recognise changes in emotional functioning, physical health, interpersonal relationships and professional behaviour.

4.1 Emotional Manifestations

Emotional symptoms may include persistent sadness, irritability, frustration, emotional exhaustion, helplessness, reduced enthusiasm and difficulty experiencing satisfaction from patient care. Nurses may become increasingly sensitive to criticism, feel overwhelmed by routine responsibilities or experience emotional detachment during patient interactions.

Some nurses may develop feelings of guilt when they cannot provide the level of care they believe patients deserve. Others may experience emotional numbness as a protective response to repeated exposure to distressing situations.

4.2 Cognitive Manifestations

Cognitive changes may involve difficulty concentrating, forgetfulness, reduced attention, indecisiveness and persistent thoughts about distressing clinical experiences. Nurses may repeatedly remember difficult patient encounters or worry excessively about possible adverse outcomes.

Persistent negative thinking may influence clinical confidence and decision-making. Nurses may begin to question their professional competence or believe that their interventions are ineffective, even when they are providing appropriate care.

4.3 Physical Manifestations

Physical symptoms may include fatigue, headaches, muscular tension, gastrointestinal discomfort, sleep disturbances and changes in appetite. Some nurses may experience persistent tiredness despite adequate rest, while others may report difficulty initiating or maintaining sleep.

When these symptoms continue, they may affect physical health, concentration and the ability to perform professional responsibilities effectively.

4.4 Behavioural Manifestations

Behavioural changes may include social withdrawal, reduced communication, increased absenteeism, irritability, avoidance of emotionally demanding patients and decreased participation in professional activities.

Some nurses may develop unhealthy coping practices, such as excessive caffeine consumption, increased alcohol use or inappropriate reliance on other substances. These behaviours may temporarily reduce distress but can create additional health problems.

4.5 Professional Manifestations

Professional manifestations may include reduced empathy, decreased job satisfaction, diminished therapeutic engagement, frequent thoughts about leaving the profession and difficulty maintaining professional boundaries.

A nurse who previously demonstrated enthusiasm for therapeutic communication may begin to avoid prolonged patient interactions or respond mechanically to emotional disclosures. Such changes should be recognised as possible indicators of occupational distress rather than automatically interpreted as a lack of professional commitment.

5. Impact of Compassion Fatigue on Mental Health Nurses

Compassion fatigue affects more than individual emotional wellbeing. It may influence professional functioning, interpersonal relationships, healthcare quality and the sustainability of mental health services.

5.1 Psychological Health of Nurses

Persistent compassion fatigue may contribute to anxiety, depressive symptoms, emotional exhaustion and secondary traumatic stress. Nurses may experience reduced self-confidence, increased psychological distress and difficulty maintaining a healthy work-life balance.

copyright©: Scientific Journal

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

Cocker and Joss, in their systematic review of healthcare and community service professionals, identified associations between compassion fatigue, burnout and secondary traumatic stress, highlighting the importance of interventions to protect workers' psychological wellbeing [2].

5.2 Quality of Patient Care

Compassion fatigue may reduce nurses' emotional availability, patience and capacity for therapeutic engagement. Patients may receive less individualised emotional support, and communication may become increasingly task-oriented.

In mental health settings, this is particularly important because therapeutic relationships contribute to patient engagement, treatment adherence, recovery and rehabilitation. Reduced empathy may weaken trust and affect the therapeutic environment.

5.3 Professional Relationships and Team Functioning

Nurses experiencing emotional exhaustion may become withdrawn, irritable or less willing to participate in teamwork. Workplace misunderstandings and interpersonal conflicts may increase, affecting communication between nursing staff and other healthcare professionals.

A supportive professional environment is therefore necessary not only for individual wellbeing but also for effective multidisciplinary care.

5.4 Absenteeism and Workforce Retention

Persistent compassion fatigue may contribute to reduced work satisfaction, sickness absence, professional disengagement and intentions to leave employment. This can create additional staffing difficulties and increase the workload of remaining nurses.

Marshman and colleagues highlighted the potential implications of compassion fatigue for mental health nursing recruitment, retention and the sustainability of compassionate care [3].

5.5 Family and Personal Relationships

The emotional effects of demanding clinical work may continue beyond working hours. Nurses may experience irritability, reduced interest in family activities, emotional withdrawal and difficulty relaxing at home.

Repeatedly thinking about patients' traumatic experiences may interfere with personal relationships and recreational activities. Maintaining boundaries between professional responsibilities and personal life is therefore an important component of prevention.

6. Assessment and Early Identification of Compassion Fatigue

Assessment is essential for identifying nurses who may be experiencing compassion fatigue and determining the level of support required. Because compassion fatigue is not a standalone psychiatric diagnosis, assessment should focus on professional quality of life, occupational stress, secondary traumatic stress symptoms and associated psychological difficulties.

6.1 Self-Assessment and Professional Reflection

Nurses should be encouraged to recognise changes in their emotional responses, physical health, interpersonal relationships and professional behaviour. Reflective self-assessment may help identify persistent exhaustion, emotional detachment, reduced empathy and difficulty recovering after demanding clinical experiences.

Regular reflection should be considered a professional wellbeing activity rather than an indication of weakness. Nurses should be encouraged to seek assistance when symptoms begin to affect personal functioning or patient care.

6.2 Professional Quality of Life Scale

The Professional Quality of Life Scale (ProQOL) developed by Stamm is one of the commonly used instruments for examining professional quality of life among individuals involved in helping professions [4].

The ProQOL-5 contains 30 items assessing three dimensions: compassion satisfaction, burnout and secondary traumatic stress. The instrument can help identify patterns associated with professional wellbeing and occupational distress.

It is important to interpret ProQOL results according to the instrument's scoring guidance. The scale is not a diagnostic tool and should not be used as the sole basis for determining an individual's psychological condition or employment suitability.

6.3 Assessment of Burnout

The Maslach Burnout Inventory is another established instrument used to assess burnout through emotional exhaustion, depersonalisation and personal accomplishment [6].

It may be useful when organisational stress and prolonged occupational exhaustion are suspected. However, burnout assessment alone does not adequately capture every aspect of secondary traumatic stress or compassion fatigue.

6.4 Assessment of Secondary Traumatic Stress

The Secondary Traumatic Stress Scale and other validated trauma-related instruments may be considered when nurses report intrusive thoughts, avoidance, heightened alertness or distress associated with indirect exposure to traumatic experiences.

Clinical assessment should also explore sleep, anxiety, depressive symptoms, substance use and functional impairment. Nurses with significant psychological symptoms should be referred for appropriate professional evaluation rather than relying exclusively on occupational screening instruments.

6.5 Organisational Assessment

copyright©: Scientific Journal

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

Nursing administrators should assess workplace conditions that may contribute to compassion fatigue. Relevant factors include staffing adequacy, patient acuity, workload distribution, frequency of exposure to traumatic incidents, workplace aggression, availability of supervision, opportunities for rest and access to psychological support.

Organisational assessment is particularly important because an individual nurse's distress may reflect systemic workplace problems rather than personal coping difficulties alone.

7. Prevention of Compassion Fatigue Through Individual Nursing Strategies

Prevention should begin with the development of healthy coping practices, emotional awareness and appropriate professional boundaries. Individual strategies can strengthen the nurse's capacity to manage demanding clinical experiences, although they must be supported by organisational interventions.

7.1 Self-Care and Personal Wellbeing

Self-care involves deliberate activities that maintain physical, psychological, emotional and social health. Mental health nurses should establish realistic routines that include adequate sleep, balanced nutrition, regular physical activity, relaxation and recreational activities.

Maintaining interests outside professional responsibilities can help create psychological distance from stressful clinical experiences. Nurses should also recognise personal limitations and avoid continuously sacrificing their health to meet workplace demands.

Self-care should not be interpreted as an individual responsibility that replaces the employer's duty to provide safe and supportive working conditions.

7.2 Mindfulness and Relaxation Techniques

Mindfulness involves paying attention to present experiences with awareness and without excessive judgement. Practices such as mindful breathing, meditation, progressive muscle relaxation and guided imagery may help nurses regulate emotional responses and manage stress.

Mindfulness can be incorporated into short breaks, structured wellbeing programmes or individual relaxation routines. Regular practice may improve emotional awareness and help nurses recognise stress responses before they become overwhelming.

7.3 Maintaining Professional Boundaries

Appropriate professional boundaries are essential for maintaining therapeutic relationships while protecting nurses' emotional wellbeing. Nurses should demonstrate empathy without assuming responsibility for every patient outcome.

Understanding the limits of professional responsibility, using appropriate communication and seeking supervision during emotionally demanding cases can reduce excessive emotional involvement.

7.4 Reflective Practice

Reflective practice allows nurses to examine clinical experiences, identify emotional responses and consider alternative approaches to challenging situations. Reflection may occur through personal journaling, structured supervision, reflective discussions or professional learning activities.

For example, following a difficult patient suicide attempt, a nurse may reflect on the emotional impact of the incident, the clinical decisions involved, the support received and the lessons that can inform future practice. Reflection should be conducted in a supportive environment and should not become a process of self-blame.

7.5 Building Psychological Resilience

Resilience refers to the ability to adapt and recover when confronted with stress, adversity or challenging experiences. Mental health nurses can strengthen resilience through problem-solving skills, emotional regulation, realistic expectations, social support and professional learning.

Resilience does not mean tolerating unsafe working conditions or remaining unaffected by traumatic experiences. Rather, it involves developing adaptive responses while recognising when additional assistance is necessary.

7.6 Social Support and Work-Life Balance

Supportive relationships with family members, friends and colleagues can reduce feelings of isolation and improve emotional recovery. Nurses should be encouraged to maintain meaningful social connections and participate in activities unrelated to professional responsibilities.

Maintaining a healthy work-life balance also requires appropriate management of working hours, leave, rest periods and personal commitments.

8. Organisational Strategies for Preventing Compassion Fatigue

Although individual strategies are valuable, compassion fatigue cannot be adequately prevented through self-care alone. Healthcare institutions have a major responsibility to create environments that protect the psychological wellbeing of mental health nurses.

8.1 Supportive Leadership and Management

Nursing leaders should recognise compassion fatigue as a legitimate occupational health concern. Managers can promote

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

supportive communication, encourage staff to discuss difficulties and respond constructively when nurses request assistance.

Leadership practices should include fair workload distribution, professional recognition, transparent decision-making and opportunities for staff participation in clinical and organisational decisions.

Marshman and colleagues identified strong leadership and positive workplace cultures as important protective considerations for compassion fatigue among mental health nurses [3].

8.2 Adequate Staffing and Workload Management

Healthcare institutions should regularly assess staffing requirements according to patient acuity, clinical risks, service demands and nursing responsibilities.

Adequate staffing can improve opportunities for therapeutic engagement, reduce excessive workload and support appropriate distribution of emotionally demanding cases.

Nurse managers should also monitor repeated overtime, missed breaks and excessive consecutive shifts because these conditions may contribute to occupational exhaustion.

8.3 Clinical Supervision

Clinical supervision provides a structured professional relationship in which nurses can discuss clinical experiences, examine therapeutic interactions, identify learning needs and receive professional guidance.

In mental health nursing, supervision is especially important because clinical work frequently involves complex emotional and interpersonal situations. It can help nurses recognise emotional reactions, maintain professional boundaries and improve therapeutic practice.

Supervision should be regular, accessible and conducted by appropriately trained professionals. It should be distinguished from routine administrative performance evaluation to ensure that nurses feel comfortable discussing difficulties.

8.4 Peer Support and Reflective Group Discussions

Peer support programmes provide opportunities for nurses to share experiences, discuss challenges and receive encouragement from colleagues.

Structured reflective groups can help nurses understand that emotional reactions to difficult clinical situations are not necessarily individual failures. Group discussions may also identify common workplace stressors requiring organisational attention.

Confidentiality and respectful communication should be maintained during these activities.

8.5 Psychological Counselling and Employee Assistance

Healthcare institutions should provide confidential psychological support services for nurses experiencing significant occupational distress.

Employee assistance programmes, counselling services, mental health consultations and referral pathways can help nurses access professional support before symptoms become severe.

Access to psychological services should be simple, confidential and free from unnecessary professional stigma.

8.6 Post-Incident Support

Following patient suicide, serious self-harm, workplace violence or other traumatic clinical events, nurses may require immediate emotional support and further assessment.

Organisations should provide appropriate post-incident support, opportunities for voluntary reflective discussion and referral to professional counselling when necessary.

Such support should avoid blame-oriented discussions and should distinguish psychological care from formal clinical incident investigation.

9. Role of Psychiatric Nurse Educators in Prevention

Nursing education is important for developing awareness of compassion fatigue before nurses enter demanding clinical environments. Psychiatric nurse educators can incorporate professional wellbeing, emotional regulation, reflective practice and occupational mental health into undergraduate and postgraduate nursing curricula.

Student nurses should be educated about the emotional demands of therapeutic relationships, professional boundaries and the importance of seeking supervision. Clinical placements should provide opportunities to discuss challenging patient interactions under appropriate faculty guidance.

Continuing nursing education programmes can address the recognition of secondary traumatic stress, burnout, compassion fatigue and professional quality of life. Workshops may include case discussions, self-assessment activities, mindfulness training, stress management and reflective exercises.

Educators should also help students understand that compassionate care does not require unlimited emotional availability. Professional empathy must be combined with self-awareness, appropriate boundaries and access to support.

10. Evidence-Based Interventions for Compassion Fatigue

Several interventions have been explored to reduce compassion fatigue and improve professional wellbeing among nurses and other healthcare workers. However, the strength of evidence varies, and no single intervention is universally effective.

10.1 Mindfulness-Based Interventions

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

Mindfulness-based interventions may improve emotional awareness, stress regulation and psychological wellbeing. They can involve meditation, mindful breathing, body awareness and structured group activities.

These programmes may be particularly helpful when nurses experience persistent emotional arousal and difficulty disengaging from stressful clinical events. Nevertheless, mindfulness should be offered as a supportive intervention rather than a substitute for addressing excessive workload or unsafe working conditions.

10.2 Resilience Training

Resilience training programmes aim to strengthen coping skills, problem-solving abilities, emotional regulation and adaptive responses to occupational stress.

Such programmes may incorporate psychological education, reflective exercises, communication skills and stress-management techniques. The objective is to improve nurses' ability to manage demanding situations without ignoring the structural causes of distress.

10.3 Compassion Fatigue Education

Educational interventions help nurses understand the meaning, risk factors, warning signs and consequences of compassion fatigue. Improved awareness may encourage early self-recognition and appropriate help-seeking.

Cocker and Joss found that intervention research targeting compassion fatigue in healthcare and community service workers was developing, with several included studies demonstrating improvements in burnout or secondary traumatic stress outcomes [2].

10.4 Compassion Satisfaction Enhancement

Prevention should also focus on strengthening the positive aspects of professional caregiving. Recognition of achievements, opportunities for professional development, positive feedback, supportive teamwork and participation in meaningful patient recovery activities may improve compassion satisfaction.

Alreshidi and Rayani examined predictors of compassion satisfaction, compassion fatigue and burnout among nursing professionals in Saudi Arabia, highlighting the relevance of professional and workplace factors to nurses' quality of life [7].

10.5 Multicomponent Interventions

Combining several strategies may be more appropriate than depending on a single intervention. A programme may integrate compassion fatigue education, mindfulness, reflective practice, peer support, clinical supervision and organisational improvements.

Garnett and colleagues' 2023 scoping review of compassion fatigue among healthcare providers identified the importance of understanding the phenomenon within broader workforce wellbeing and organisational contexts [8].

The 2022 systematic review by Marshman and colleagues also supports an integrated approach involving awareness, individual resilience, clinical supervision and supportive workplace cultures for mental health nurses [3].

11. Developing a Compassion Fatigue Prevention Programme in Psychiatric Settings

Healthcare institutions can establish structured prevention programmes to integrate professional wellbeing into routine nursing administration. Such programmes should be designed according to the needs of individual psychiatric units and should include both preventive and supportive components.

The first step involves assessing the existing workplace environment, including staffing, workload, exposure to traumatic events, workplace violence and access to supervision. This assessment helps identify specific occupational risks.

The second step involves educating nursing staff about compassion fatigue, burnout, secondary traumatic stress and compassion satisfaction. Educational sessions should include early warning signs, self-assessment and available support services.

The third step involves establishing regular clinical supervision, peer-support meetings and opportunities for reflective practice. Nurses should be encouraged to discuss difficult patient interactions in a psychologically safe environment.

The fourth step involves implementing individual wellbeing initiatives, such as mindfulness sessions, relaxation activities, physical activity programmes and stress-management workshops. These should be voluntary, accessible and adapted to shift-based working arrangements.

The fifth step involves creating clear referral pathways for nurses who require professional psychological assistance. Confidentiality must be maintained, and staff should not fear discrimination or professional punishment for seeking help.

Finally, programme effectiveness should be evaluated through staff feedback, professional quality-of-life assessments, absenteeism, retention, participation rates and workplace climate indicators.

12. Challenges in the Prevention of Compassion Fatigue

Several barriers may affect the implementation of effective prevention strategies in psychiatric nursing settings.

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

Stigma surrounding psychological support: Some nurses may consider seeking psychological assistance a sign of professional weakness. This belief can delay early identification and intervention. Nursing leaders should promote a culture in which help-seeking is recognised as responsible professional behaviour.

Inadequate institutional resources: Limited staffing, financial constraints and insufficient access to trained supervisors may restrict the implementation of prevention programmes. Institutions should consider professional wellbeing an essential component of workforce development rather than an optional activity.

Limited awareness and training: Nurses and managers may have difficulty distinguishing compassion fatigue from burnout, general stress and secondary traumatic stress. Structured educational programmes can improve understanding and encourage appropriate assessment.

Workload and time constraints: Nurses working in busy psychiatric units may have limited opportunities to attend counselling sessions, reflective discussions or wellbeing activities. Programmes should therefore be incorporated into working schedules.

Overemphasis on individual responsibility: Prevention initiatives may focus excessively on resilience and self-care while ignoring staffing shortages, workplace aggression and inadequate management support. Such approaches may unintentionally place responsibility for organisational problems on individual nurses.

Insufficient research in mental health nursing: Although research has increased, there remains a need for larger longitudinal studies examining the experiences of mental health nurses across different clinical settings and cultural contexts. Marshman and colleagues specifically highlighted limitations in the existing quantitative literature and the need to understand mental health nurses' lived experiences of compassion fatigue [3].

13. Future Directions for Nursing Research and Practice

Future research should investigate the long-term effects of compassion fatigue on psychiatric nursing practice, patient outcomes, professional retention and healthcare quality. Longitudinal studies may help identify how occupational exposure, personal coping resources and organisational conditions interact over time.

Research should also examine the effectiveness of structured clinical supervision, peer-support interventions, mindfulness-based programmes and organisational prevention strategies. Comparative studies could identify which combinations of interventions provide sustainable benefits in different psychiatric settings.

Greater attention should be given to nurses working in community mental health services, addiction treatment facilities, forensic psychiatric units, child and adolescent mental health services and crisis intervention settings. These areas may involve distinct emotional demands that are not adequately represented in general nursing research.

In India and other low- and middle-income countries, additional research is needed to examine the influence of nurse staffing, workload, professional recognition, cultural expectations and access to psychological support. Locally relevant interventions should consider institutional resources and the specific occupational challenges faced by psychiatric nurses.

Digital mental health resources may also provide opportunities for confidential wellbeing education, guided relaxation, self-assessment and access to counselling. However, digital programmes should complement professional supervision and institutional support rather than replace direct human interaction.

Ultimately, compassion fatigue prevention should become an integral part of psychiatric nursing education, occupational health policies, clinical governance and healthcare workforce planning.

14. Conclusion

Compassion fatigue is an important occupational concern in mental health nursing because of the continuous emotional demands associated with caring for individuals experiencing psychological distress, trauma, suicidal behaviour and complex psychosocial difficulties. Repeated exposure to suffering, workplace violence, heavy workloads, ethical dilemmas and inadequate organisational support may gradually affect nurses' emotional wellbeing, physical health, professional relationships and capacity to provide compassionate care.

Early recognition requires attention to emotional, cognitive, physical, behavioural and professional manifestations. Instruments such as the Professional Quality of Life Scale can support the assessment of professional wellbeing, while clinical supervision and reflective practice provide opportunities to recognise and address occupational distress.

Prevention requires a comprehensive approach involving individual self-care, mindfulness, resilience development, appropriate professional boundaries, peer support, clinical supervision, supportive leadership and adequate organisational resources. Importantly, compassion fatigue should not be viewed simply as an individual's failure to cope. Healthcare institutions must recognise their responsibility to establish psychologically safe working environments and accessible support systems.

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

Psychiatric nurses are essential contributors to recovery-oriented mental healthcare. Protecting their emotional wellbeing is therefore not only an occupational health priority but also an important component of patient safety, therapeutic effectiveness and healthcare quality. Building a professional culture that values both compassionate patient care and compassionate care for nurses is essential for sustaining the mental health nursing workforce.

References

1. Figley CR, editor. **Compassion Fatigue: Coping with Secondary Traumatic Stress Disorder in Those Who Treat the Traumatized**. New York: Brunner/Mazel; 1995.
2. Cocker F, Joss N. Compassion fatigue among healthcare, emergency and community service workers: a systematic review. Int J Environ Res Public Health. 2016;13(6):618. doi:10.3390/ijerph13060618.
3. Marshman C, Hansen A, Munro I. Compassion fatigue in mental health nurses: a systematic review. J Psychiatr Ment Health Nurs. 2022;29(4):529-543. doi:10.1111/jpm.12812.
4. Stamm BH. The Concise ProQOL Manual. 2nd ed. Pocatello, ID: ProQOL.org; 2010.
5. World Health Organization. **International Classification of Diseases for Mortality and Morbidity Statistics (11th Revision)**. Geneva: World Health Organization; 2019.
6. Maslach C, Schaufeli WB, Leiter MP. Job burnout. Annu Rev Psychol. 2001;52:397-422. doi:10.1146/annurev.psych.52.1.397.
7. Alreshidi SM, Rayani AM. Predictors of compassion satisfaction, compassion fatigue, and burnout among nursing professionals in a medical city in Saudi Arabia. Risk Manag Healthc Policy. 2023;16:2883-2892. doi:10.2147/RMHP.S430082.
8. Garnett A, Hui L, Oleynikov C, Boamah S. Compassion fatigue in healthcare providers: a scoping review. BMC Health Serv Res. 2023;23(1):1336. doi:10.1186/s12913-023-10356-3.
9. Alreshidi SM. Compassion fatigue prevalence and risk factors among Saudi psychiatric nurses: a cross-sectional study. Medicine (Baltimore). 2023;102(45):e35975. doi:10.1097/MD.0000000000035975.
10. Figley CR. Compassion fatigue: psychotherapists' chronic lack of self care. J Clin Psychol. 2002;58(11):1433-1441.
11. Joinson C. Coping with compassion fatigue. Nursing. 1992;22(4):116-121.
12. Hunsaker S, Chen HC, Maughan D, Heaston S. Factors that influence the development of compassion fatigue, burnout, and compassion satisfaction in emergency department nurses. J Nurs Scholarsh. 2015;47(2):186-194. doi:10.1111/jnu.12122.
13. Duarte J, Pinto-Gouveia J. Empathy and feelings of guilt experienced by nurses: a cross-sectional study of their role in burnout and compassion fatigue symptoms. Appl Nurs Res. 2017;35:42-47.
14. Potter P, Deshields T, Berger J, Clarke M, Olsen S, Chen L. Evaluation of a compassion fatigue resiliency program for oncology nurses. Oncol Nurs Forum. 2013;40(2):180-187.
15. Mealer M, Jones J, Newman J, McFann KK, Rothbaum B, Moss M. The presence of resilience is associated with a healthier psychological profile in intensive care unit nurses: results of a national survey. Int J Nurs Stud. 2012;49(3):292-299.
16. Rushton CH, Batcheller J, Schroeder K, Donohue P. Burnout and resilience among nurses practicing in high-intensity settings. Am J Crit Care. 2015;24(5):412-420.
17. Figley CR. Treating compassion fatigue. New York: Brunner-Routledge; 2002.
18. Beck CT. Secondary traumatic stress in nurses: a systematic review. Arch Psychiatr Nurs. 2011;25(1):1-10.
19. Hunsaker S, Chen HC, Maughan D, Heaston S. Factors influencing compassion fatigue, burnout, and compassion satisfaction in emergency nurses. J Nurs Scholarsh. 2015;47(2):186-194.
20. Coetzee SK, Klopper HC. Compassion fatigue within nursing practice: a concept analysis. Nurs Health Sci. 2010;12(2):235-243. doi:10.1111/j.1442-2018.2010.00526.x.
21. Sinclair S, Raffin-Bouchal S, Venturato L, Mijovic-Kondejewski J, Smith-MacDonald L. Compassion fatigue: a meta-narrative review of the healthcare literature. Int J Nurs Stud. 2017;69:9-24.
22. Stamm BH. Measuring compassion satisfaction as well as fatigue: development history of the Compassion Satisfaction and Fatigue Test. In: Figley CR, editor. Treating Compassion Fatigue. New York: Brunner-Routledge; 2002.
23. Maslach C, Leiter MP. Understanding the burnout experience: recent research and its implications for psychiatry. World Psychiatry. 2016;15(2):103-111. doi:10.1002/wps.20311.
24. Figley CR. Compassion fatigue: an introduction. In: Figley CR, editor. Compassion Fatigue: Coping with Secondary Traumatic Stress Disorder in Those Who Treat the Traumatized. New York: Brunner/Mazel; 1995. p. 1-20.
25. World Health Organization. **Guidelines on Mental Health at Work**. Geneva: World Health Organization; 2022.
26. International Council of Nurses. **The ICN Code of Ethics for Nurses**. Geneva: International Council of Nurses; 2021.
27. National Academies of Sciences, Engineering, and Medicine. **Taking Action Against Clinician Burnout: A Systems**

SCIENTIFIC JOURNAL
www.scientificjournal.in
JOURNAL PUBLICATIONS INDEXED IN
www.scientificjournal.in YEAR: 2025 VOLUME: 3 ISSUE: 2 ISSN: 3107-4162

Approach to Professional Well-Being. Washington, DC: National Academies Press; 2019.
28. Sabo BM. Reflecting on the concept of compassion fatigue. Online J Issues Nurs. 2011;16(1):1.
29. Sorenson C, Bolick B, Wright K, Hamilton R. Understanding compassion fatigue in healthcare providers: a review of current literature. J Nurs Scholarsh. 2016;48(5):456-465.
30. Ledoux K. Understanding compassion fatigue: understanding compassion. J Adv Nurs. 2015;71(9):2041-2050.

copyright©: Scientific Journal